

## Lattice Polygons in the Plane and the Number 12

JOHN M. BURNS AND DAVID O'KEEFFE

### 1. INTRODUCTION

A convex polygon  $\mathcal{P}$  in  $\mathbb{R}^2$  all of whose vertices have integer coordinates is called a convex lattice polygon. If the polygon has  $n$  lattice points on its boundary, represented by the vectors  $p_1, \dots, p_n$  (in anticlockwise order), then we say that the length  $l(\mathcal{P})$  is  $n$ . The dual (convex) lattice polygon  $\mathcal{P}^\vee$  is by definition the convex hull of the difference vectors  $q_i = p_{i+1} - p_i$ , where indices throughout the note are considered modulo  $n$ . In this note we will give a simple proof of the fact that if  $\mathcal{P}$  be a convex lattice polygon in  $\mathbb{R}^2$  whose only interior lattice point is the origin, then  $l(\mathcal{P}) + l(\mathcal{P}^\vee) = 12$ . This result has its origin in a correspondence between convex lattice polygons and certain toric varieties, and has several ingenious proofs (see [1] and [2]). These proofs use either Noether's formula or modular forms to explain the occurrence of the number 12. Our proof observes that if  $l(\mathcal{P})$  increases by one then  $l(\mathcal{P}^\vee)$  decreases by one, so that their sum is constant. The number 12 then appears as 3 (the smallest possible length) plus 9 (the largest possible length.) Our proof (which grew out of the second authors project, supervised by the first author) also supports the suggestion in [2] that the number 3 can be viewed as a discrete analogue of  $\pi$  in this context.

### 2. THE THEOREM

**Theorem 2.1.** *Let  $\mathcal{P}$  be a convex lattice polygon in  $\mathbb{R}^2$  whose only interior lattice point is the origin, and let  $\mathcal{P}^\vee$  be its dual. Then  $l(\mathcal{P}) + l(\mathcal{P}^\vee) = 12$ .*

It is not hard to see that up to the action of  $GL(2, \mathbb{Z})$  there are 16 such polygons but we will not need this fact. We will however use the following result of P. Scott [3], the proof of which relies only on elementary geometry.

**Lemma 2.1.** *Let  $\mathcal{P}$  be a convex lattice polygon in the plane with  $l(\mathcal{P})$  boundary lattice points and  $c \geq 1$  interior points then  $l(\mathcal{P}) \leq 2c + 7$ .*

We will also use picks formula, which in our context states:

**Lemma 2.2.** *The area of a lattice polygon  $\mathcal{P}$  is given by the formula  $A(\mathcal{P}) = c + l(\mathcal{P})/2 - 1$ .*

In our case  $c = 1$ , so that  $l(\mathcal{P}) \leq 9$  and  $A(\mathcal{P}) = l(\mathcal{P})/2$ . Since we will think of the plane as the complex plane  $\mathbb{C}$ , we will label the vectors  $p_i$  and  $q_i$  as  $z_i$  and  $w_i$  respectively and we let  $\gamma_i$  denote the straight line path (parameterised on  $[0, 1]$ ) joining  $z_i$  to  $z_{i+1}$ . We will fix  $z_1 = 1$  throughout unless we state otherwise. We recall that for a piecewise smooth curve  $\gamma$  in  $\mathbb{C}$  we have  $\int_{\gamma} \bar{z} dz = 2iA$  where  $A$  denotes the area enclosed by  $\gamma$ . In particular a convex lattice polygon with vertices  $z_1, \dots, z_n$  has area

$$\begin{aligned} & -\frac{i}{2} \sum_{i=1}^n \int_0^1 (tz_{i+1} + (1-t)z_i)(z_{i+1} - z_i) dt \\ &= -\frac{i}{4} \sum_{i=1}^n [|z_{i+1}|^2 - |z_i|^2 + \bar{z}_i z_{i+1} - z_i \bar{z}_{i+1}] \\ &= -\frac{i}{4} \sum_{i=1}^n [\bar{z}_i z_{i+1} - z_i \bar{z}_{i+1}]. \end{aligned}$$

It will be convenient to introduce the notation  $A_{ij} = -\frac{i}{4}(\bar{z}_i z_j - z_i \bar{z}_j)$ , namely the signed area of the oriented triangle with vertices  $o, z_i, z_j$ . Similarly, we let  $A_{ij}^{\vee} = -\frac{i}{4}(\bar{w}_i w_j - w_i \bar{w}_j)$ , so that

$$A_{ii+1}^{\vee} = -\frac{i}{4}(\bar{z}_i z_{i+1} - z_i \bar{z}_{i+1} + \bar{z}_{i+1} z_{i+} - z_{i+1} \bar{z}_{i+2} + \bar{z}_{i+2} z_i - z_i \bar{z}_{i+2}).$$

In summary  $A_{ii+1}^{\vee} = A_{ii+2} + A_{i+1i+2} + A_{i+2i}$ , so that  $l(\mathcal{P}^{\vee}) = 2l(\mathcal{P}) - \sum_{i=1}^n A_{ii+2}$ . This immediately yields that when  $l(\mathcal{P}) = n = 3$  we have  $l(\mathcal{P}^{\vee}) = A_{12}^{\vee} + A_{23}^{\vee} + A_{31}^{\vee} = (A_{12} + A_{23} + A_{31}) + (A_{23} + A_{31} + A_{12}) + (A_{31} + A_{12} + A_{23}) = 3(A_{31} + A_{12} + A_{23}) = 3 \times 3 = 9$  (since  $A_{ii+1} = 1$  for all  $i$ ). If  $l(\mathcal{P}) = n = 4$ ,  $\sum_{i=1}^n A_{ii+2} = A_{13} + A_{24} + A_{31} + A_{42} = 0$  (since  $A_{ij} = -A_{ji} \forall i, j$ ) and we have  $l(\mathcal{P}^{\vee}) = 2l(\mathcal{P}) = 8$ . In future we will denote the sum  $\sum_{i=1}^n A_{ii+2}$  by  $d(\mathcal{P})$ , and show that it increases by 3 when  $l(\mathcal{P})$  increases by 1, thus keeping  $l(\mathcal{P}) + l(\mathcal{P}^{\vee})$  constant.

*Proof.* The case  $l(\mathcal{P}) = n = 5$  is the crucial case as all others essentially follow from this one. Here since  $A(\mathcal{P}) = 5/2$  and no edge

contains more than 3 vertices, we may centre  $\mathcal{P}$  at the origin with its vertices on the unit square. Since one of the 4 lines  $x = 0$ ,  $y = 0$ ,  $y = \pm x$  must contain 2 of the 5 vertices of  $\mathcal{P}$ , we will assume that it is the  $x$ -axis. We may also assume that 2 ( $z_2$  and  $z_3$ ) of the remaining 3 vertices lie above the  $x$ -axis and that the remaining vertex  $z_5$  lies below it. It is immediate that  $A_{41} = 0$  ( $z_4, o$ , and  $z_1$  being collinear). We first consider the case where  $\mathcal{P}$  has an edge of length 2 (i.e., containing 3 vertices). This edge consists either of the points  $\{z_5, z_1, z_2\}$  or  $\{z_3, z_4, z_5\}$ , so that either  $A_{52} = 2$  or  $A_{35} = 2$ . In the former case  $A_{35} = -1$ , and in the latter  $A_{52} = -1$ . In both cases the remaining  $A_{ii+2}$  are equal to 1, so that  $\sum_{i=1}^n A_{ii+2} = 3$  and we are done. When  $\mathcal{P}$  has no edge of length 2, we must have an additional  $A_{ii+2} = 0$  on account of the fact that  $z_5$  doesn't lie on a vertical edge of length 2. This forces the remaining  $A_{ii+2}$  to be 1, and again  $\sum_{i=1}^n A_{ii+2} = 3$ . When  $l(\mathcal{P}) = 6$ , we can (after relabelling if necessary) delete the vertex  $z_6$  from  $\mathcal{P}$  to obtain a convex lattice polygon  $\mathcal{P}'$  containing the origin, and  $d(\mathcal{P}) - d(\mathcal{P}') = A_{13} + A_{35} + A_{51} + A_{24} + A_{46} + A_{62} - (A_{13} + A_{35} + A_{52} + A_{24} + A_{41}) = A_{51} + A_{46} + A_{62} + A_{25} + A_{14} = 3$ . The last equality follows from the observation that  $A_{51} + A_{46} + A_{62} + A_{25} + A_{14} = d(\mathcal{P}'')$  where  $\mathcal{P}''$  is either the convex lattice pentagon  $\{z_1, z_2, z_4, z_5, z_6\}$  containing the origin, so that  $d(\mathcal{P}'') = 3$  by above, or else the convex lattice hexagon (the above pentagon with the origin adjoined) with no interior lattice point, where the computation is trivial. We now have  $l(\mathcal{P}^\vee) = 2l(\mathcal{P}) - (d(\mathcal{P}') + 3) = 12 - 6 = 6$ . It is intriguing that the cases  $l(\mathcal{P}) = 7, 8, 9$  are identical to that of  $l(\mathcal{P}) = 6$ . In each case, just as above  $d(\mathcal{P}) - d(\mathcal{P}') = d(\mathcal{P}'')$  where  $\mathcal{P}''$  is either a convex lattice pentagon containing the origin, or the convex lattice hexagon with no interior lattice point, so that  $d(\mathcal{P}) - d(\mathcal{P}') = 3$ .  $\square$

*Remark.* Finally we point out some connections with [2]. There it is shown that the vectors  $p_i$  and  $p_{i+1}$  form a basis for the lattice  $\mathbb{Z}^2$  with the same orientation as the standard basis  $\{(1, 0), (0, 1)\}$ , and that there are matrices  $M_i = \begin{pmatrix} 0 & 1 \\ -1 & d_i \end{pmatrix}$  in  $SL(2, \mathbb{Z})$  such that  $M_i \begin{pmatrix} p_{i-1} \\ p_i \end{pmatrix} = \begin{pmatrix} p_i \\ p_{i+1} \end{pmatrix}$ , where the  $2 \times 2$  matrices  $\begin{pmatrix} p_i \\ p_{i+1} \end{pmatrix}$  have the row vectors  $p_i$  and  $p_{i+1}$  as their rows. In addition we have that

$$\begin{pmatrix} 0 & 1 \\ -1 & d_n \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & d_{n-1} \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ -1 & d_1 \end{pmatrix} = I,$$

the identity matrix. It is a simple matter to see that in fact  $d_i = A_{ii+2}$  and that the above equations therefore contain some of our observations above. When  $n = 3$  they imply that  $A_{13} = A_{32} = A_{21} = -1$ . When  $n = 4$  they imply that  $A_{ii+2} = -A_{i+2i} \forall i$ . When  $n = 5$  they imply that the multi-set  $\{A_{13}, A_{35}, A_{52}, A_{24}, A_{41}\}$  is either the multi-set  $\{-1, 2, 1, 0, 1\}$  or else  $\{0, 1, 1, 0, 1\}$ . In either case  $d(\mathcal{P}) = \sum_{i=1}^n A_{ii+2} = 3$ . We also observe that our proof supports the suggestion in [2] that 3 can be viewed as a discrete analogue of  $\pi$ . Clearly  $d(\mathcal{P})$  has an interpretation as the sum of the “discrete exterior angles” of  $\mathcal{P}$  (as defined in [2].) We have shown that increasing  $l(\mathcal{P})$  by one increases this sum by 3, whereas the sum of the exterior angles increases by  $\pi$  for a general polygon.

#### REFERENCES

- [1] W. Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, No. 82, Princeton University Press 1993.
- [2] B. Poonen and F. Rodriguez-Villegas, *Lattice polygons and the number 12*, Amer. Math. Monthly **107** (2000), 238–250.
- [3] P. Scott, *On convex lattice polygons*, Bull. Austral. Math. Soc. **15** (1976), 395–399.

John M. Burns and David O’Keeffe,  
 Department of Mathematics  
 National University of Ireland,  
 Galway, Ireland  
 john.burns@nuigalway.ie

*Received on 25 October 2005.*